

Dialog DataStar

options

logoff

feedback

help

databases

easy
search**Advanced Search:**

Inspec - 1898 to date (INZZ)

limit

Search history:

No.	Database	Search term	Info added since	Results	
CP		[Clipboard]		0	-
1	INZZ	diamondoid OR adamant\$	unrestricted	778	show titles
3	INZZ	amant\$	unrestricted	211	show titles
4	INZZ	3 OR 1	unrestricted	987	show titles
5	INZZ	4 AND (luminophor\$ OR luminesc\$ OR fluorophor\$ OR fluoresc\$ OR photophor\$ OR photoluminesc\$ OR photofluoresc\$ OR photon OR chemiluminesc\$ OR triboluminesc\$ OR photomultiplier OR ccd OR charge\$1 ADJ coupl\$3 OR spectra OR spectrum OR spectrophotomet\$ OR optical)	unrestricted	390	show titles
6	INZZ	(luminophor\$ OR luminesc\$ OR fluorophor\$ OR fluoresc\$ OR photophor\$ OR photoluminesc\$ OR photofluoresc\$ OR photon OR chemiluminesc\$ OR triboluminesc\$ OR photomultiplier OR ccd OR charge\$1 ADJ coupl\$3 OR spectra OR spectrum OR spectrophotomet\$ OR optical) SAME (diamondoid OR adamant\$ OR amant\$)	unrestricted	249	show titles
7	INZZ	(luminophor\$ OR luminesc\$ OR fluorophor\$ OR fluoresc\$ OR photophor\$ OR photoluminesc\$ OR photofluoresc\$ OR photon OR chemiluminesc\$ OR triboluminesc\$ OR photomultiplier OR ccd OR charge\$1 ADJ coupl\$3 OR spectrophotomet\$ OR optical) SAME (diamondoid OR adamant\$ OR amant\$)	unrestricted	65	show titles

[show all](#) | [hide](#) | [delete all search steps...](#) | [delete individual search steps...](#)
Enter your search term(s): [Search tips](#) ☐ Thesaurus mapping

whole document



53 3932



options

logoff

feedback

help

databases

search
page

titles

Document

Select the documents you wish to save or order by clicking the box next to the document, or click the link above the document to order directly.

previous
documentsnext
documents

save

locally as: PDF document

search strategy: do not include the search strategy

order

copy to
clipboard
☒ Select All

[17 An integrated optical Mach-Zehnder interferometer functionalized by beta-Mac](#)
[24 Chemiluminescence of praseodymium \(III\), neodymium \(III\) and ytterbium \(III\)](#)
[29 Growth method, optical properties, and application of organic nonlinear](#)
[31 Branched quantum-chain processes in the decay of 1,2-dioxetane with participa](#)
[52 Optical activity due to deuterium substitution in adamantanone and adamant](#)
[59 Free radicals in an adamantane matrix. IX. Optical excitation and emissio](#)

Full text available at



SCIENCE @ DIRECT

USPTO Full Text Retrieval Options

open url

☒ document 17 of 65 [Order Document](#)

Inspec - 1898 to date (INZZ)

Accession number & update

0007118849 20010101.

Title

An integrated **optical** Mach-Zehnder interferometer functionalized by beta-Mach-Zehnder-cyclodextrin to monitor binding reactions.

Source

Sensors and Actuators B (Chemical), {Sens-Actuators-B-Chem-Switzerland }, 20 Nov. 2001, vol. B80, no. 2, p. 116-24, 40 refs, CODEN: SABCEB, ISSN: 0925-4005.
 Publisher: Elsevier, Switzerland.

Author(s)

Busse-S, DePaoli-M, Wenz-G, Mittler-S.

Author affiliation

Busse, S., Max-Planck-Inst. fur Polymerforschung, Mainz, Germany.

Abstract

We demonstrate an integrated **optical** Mach-Zehnder interferometer for the monitoring of guest-host reactions on immobilized beta-cyclodextrin (beta-CD). **Adamantanecarboxylic acid**, 4-tert-butylbenzoic acid and methyl orange were investigated as guest molecules. The binding constants of the three guests into the two-dimensionally arranged immobilized cyclodextrins could be determined and are a quarter to a third smaller than the values measured in the three-dimensional environment of a solution.

Descriptors

~~BIOSENSORS~~; ~~CHEMICAL-REACTIONS~~; ~~CHEMICAL-VARIABLES-MEASUREMENT~~;
~~INTEGRATED-OPTICS~~; ~~MACH-ZEHNDER-INTERFEROMETERS~~; ~~MOLECULAR-BIOPHYSICS~~;

ORGANIC-COMPOUNDS; SPECTROCHEMICAL-ANALYSIS.

Classification codes

A8780B Biosensors*;
A0760L Optical-interferometry;
A4282 Integrated-optics;
A8230 Specific-chemical-reactions-reaction-mechanisms;
A8280D Electromagnetic-radiation-spectrometry-chemical-analysis;
A8715 Molecular-biophysics;
B7230J Biosensors*;
B4140 Integrated-optics;
B7320T Chemical-variables-measurement.

Keywords

integrated-optical-Mach-Zehnder-interferometer; binding-reactions- monitoring; guest-host-reactions; 2D-environment; immobilized-beta-cyclodextrin; **Adamantanecarboxylic-acid**; 4-tert-butylbenzoic-acid; methyl-orange; guest-molecules.

Treatment codes

P Practical;
X Experimental.

Language

English.

Publication type

Journal-paper.

Availability

SICI: 0925-4005(20011120)B80:2L.116:IOMZ; 1-U.
CCCC: 0925-4005/01/\$20.00.
Publisher identity number: S0925-4005(01)00889-9.

Publication year

2001.

Publication date

20011120.

Edition

2001049.

Copyright statement

Copyright 2001 IEE.

(c) 2007 The Institution of Engineering and Technology

Full text available at



SCIENCE @ DIRECT

USPTO Full Text Retrieval Options

open url

☒ **document 24 of 65** Order Document

Inspec - 1898 to date (INZZ)

Accession number & update

0006766816 20051201.

Title

Chemiluminescence of praseodymium (III), neodymium (III) and ytterbium (III) beta-diketonates in solution excited from 1,2-dioxetane decomposition and singlet-singlet energy transfer from ketone to rare-earth beta-diketonates.

Source

Journal of **Luminescence**, {J-Lumin-Netherlands}, Sept. 2000, vol. 91, no. 1-2, p. 49-58, 45 refs, CODEN: JLUMA8, ISSN: 0022-2313.
Publisher: Elsevier, Netherlands.

Author(s)

Voloshin-A-I, Shavaleev-N-M, Kazakov-V-P.

Author affiliation

Voloshin, A.I., Shavaleev, N.M., Kazakov, V.P., Inst. of Org. Chem., Ufa Sci. Center of the Russian Acad. of Sci., Oktyabrya, Russia.

Abstract

This work is concerned with the **chemiluminescence** (CL) of Nd/sup 3+/, Yb/sup 3+/, and Pr/sup 3+/-beta-diketonates in solution. **Chemiluminescent** reaction of **adamantylideneadamantane-1,2-dioxetane** (AAD) decomposition generating singlet (Ad=O/sub S/*) and triplet (Ad=O/sub T/*) excited **adamantanone** was used as a source of excited species. AAD **chemiluminescence** due to emission from Ad=O/sub S/* is quenched by Ln/sup 3+/-beta-diketonates: (a) by intermolecular singlet-singlet (S-S) energy transfer from Ad=O/sub S/* to beta-diketonate ligand levels of Ln (TTA)/sub 3/centerdot2H/sub 2/formation between AAD and Pr(FOD)/sub 3/ or Pr(DPM)/sub 3/. Corresponding Stern-Volmer quenching constants or stability constants of the complex were measured. **Chemiluminescence** spectra of Ln/sup 3+/-beta-diketonates were recorded and relative **luminescence** quantum yields compared. Yb/sup 3+/-chelates show higher **luminescence** yields compared to Nd/sup 3+/, due to a different efficiency of non-radiative energy degradation. Chemiexcitation of Ln/sup 3+/-ions in the systems studied occurs by: (a) intermolecular singlet-singlet energy transfer: Ad=O/sub S/*rightward arrowL/sub S/*rightward arrowL/sub T/*rightward arrowLn/sup 3+ and triplet excited states of the beta-diketonate ligand); (b) intermolecular triplet-triplet energy transfer: Ad=O/sub T/*rightward arrowLfrom the decomposition of AAD in the complex with Ln/sup 3+/-beta-diketonate. Efficiency of chemiexcitation pathways is different for each Ln/sup 3+/-beta-diketonate and Ln/sup 3+/-ion.

Descriptors

☒ CHEMILUMINESCENCE; ☒ NEODYMIUM-COMPOUNDS; ☒ ORGANIC-COMPOUNDS;
☒ PRASEODYMIUM-COMPOUNDS; ☒ YTTERBIUM-COMPOUNDS.

Classification codes

A7860P Chemiluminescence-condensed-matter*;
A8240T Chemiluminescence-and-chemical-laser-kinetics.

Keywords

chemiluminescence; beta-diketonates; decomposition; intermolecular- energy-transfer; lanthanide-ions; rare-earths; praseodymium; neodymium; ytterbium; singlet-singlet-energy-transfer; triplet-triplet-energy-transfer; intracomplex-energy-transfer; **luminescence**- quantum-yield; chemiexcitation; complex-formation; ketones.

Treatment codes

☒ Experimental.

Language

English.

Publication type

Journal-paper.

Availability

SICI: 0022-2313(200009)91:1/2L.49:CPNY; 1-X.
CCCC: 0022-2313/2000/\$20.00.
Publisher identity number: S0022-2313(00)00199-X.

Publication year

2000.

Publication date

20000900.

Edition

2000046.

Copyright statement

Copyright 2000 FIZ Karlsruhe.

(c) 2007 The Institution of Engineering and Technology

Full text available at



The International Society
for Optical Engineering

USPTO Full Text Retrieval Options

openurl

☒ document 29 of 65 [Order Document](#)

Inspec - 1898 to date (INZZ)

Accession number & update

0005837695 20051201.

TitleGrowth method, **optical** properties, and application of organic nonlinear **optical** crystal **2-adamantylamino-5-nitropyridine**.**Conference information**Nonlinear **Optical** Properties of Organic Materials X, San Diego, CA, USA, 30 July-1 Aug. 1997.
Sponsor(s): SPIE.**Source**Proceedings of the SPIE - The International Society for **Optical** Engineering, {Proc-SPIE-Int-Soc-Opt-Eng-USA}, 1997, vol. 3147, p. 20-9, 20 refs, CODEN: PSISDG, ISSN: 0277-786X.
Publisher: SPIE-Int. Soc. Opt. Eng, USA.**Author(s)**Yokoo-A, Yokohama-I, Takara-H, Kaino-T.**Author affiliation**

Yokoo, A., Yokohama, I., Takara, H., NTT Opto-Electron. Labs., Kanagawa, Japan.

Abstract

Organic second-order nonlinear **optical** crystals are expected to be more useful than inorganic crystals such as LiNbO_3 or KTiOPO . A long interaction length in the phase-matched direction is required to obtain devices with high wavelength conversion efficiency. We have devised an indirect laser-heated pedestal growth (ILHPG) method in which the growth direction of organic crystal can be controlled. We have successfully applied the method to grow organic nonlinear **optical** crystal **2-adamantylamino-5-nitropyridine** (AANP). The grown AANP has been characterized in terms of its linear and nonlinear **optical** properties. The results show that type II angle-tuned phase-matched wavelength conversions, such as second harmonic generation (SHG) and **optical** parametric generation (OPG), are possible in the wavelength region from 1.2 to 17 μm . This means, it is possible for AANP to provide type II angle-tuned phase-matched wavelength conversion with high efficiency for **optical** communications systems. As an example, the AANP was applied for **optical** sampling measurements. In addition, reducing absorption in the AANP crystal by using a deuterated derivative of AANP increases the SHG efficiency at 1.55 μm .

Descriptors

CRYSTAL-GROWTH-FROM-MELT; NONLINEAR-OPTICAL-SUSCEPTIBILITY; OPTICAL-COMMUNICATION-EQUIPMENT; OPTICAL-FABRICATION; OPTICAL-HARMONIC-GENERATION; OPTICAL-MATERIALS; OPTICAL-PARAMETRIC-DEVICES; WAVELENGTH-DIVISION-MULTIPLEXING.

Classification codes

A4265K Optical-harmonic-generation-frequency-conversion-parametric-oscillation-and-amplification*;
A4270F Other-optical-materials;
A8110F Crystal-growth-from-melt;
A4285D Optical-fabrication-surface-grinding;
B4340 Nonlinear-optics-and-devices*;
B4110 Optical-materials;
B0510 Crystal-growth.

Keywords

growth-method; **optical-properties**; **organic-nonlinear-optical-crystal**; **organic-second-order-nonlinear-optical-crystals**; nonlinear- susceptibility; long-interaction-length; phase-matched-direction; wavelength-conversion-efficiency; indirect-laser-heated-pedestal-growth; **nonlinear-optical-properties**; **linear-optical-properties**; type-II-angle-tuned-phase-matched-wavelength-conversion; second-harmonic-generation; **optical-parametric-generation**; **optical-communications- systems**; **optical-sampling-measurements**; deuterated-derivative; SHG-efficiency; 1.2-to-1.7- μm ; 1.55- μm .

Treatment codesX Experimental.**Numerical indexing**wavelength: 1.2E-06 to 1.7E-06 m.
wavelength: 1.55E-06 m.**Language**

English.

Publication typeConference-proceedings; Journal-paper.**Availability**

SICI: 0277-786X(1997)3147L:20:GMOP; 1-R.
CCCC: 0277-786X/97/\$10.00.

Publication year

1997.

Publication date

19970000.

Edition

1998008.

Copyright statement

Copyright 1998 IEE.

(c) 2007 The Institution of Engineering and Technology

Full text available at **USPTO Full Text Retrieval Options** [open url](#)

☒ **document 31 of 65** [Order Document](#)

Inspec - 1898 to date (INZZ)**Accession number & update**

0005796614 20051201.

Title

Branched quantum-chain processes in the decay of 1,2-dioxetane with participation of Eu(fod)/sub 3/ at the surface of a sorbent and in solution.

Source

Optics and Spectroscopy, {Opt-Spectrosc-Russia}, Nov. 1997, vol. 83, no. 5, p. 797-801, CODEN: OPSUA3, ISSN: 0030-400X.

Publisher: MAIK Nauka/Interperiodica Publishing, Russia.

Translation from: Optika i Spektroskopiya, {Opt-Spektrosk-Russia}, Nov. 1997, vol. 83, no. 5, p. 860-4, CODEN: OSFMA3, ISSN: 0030-4034.

Country of publication: Russia.

Author(s)

Kazakov-V-P, Voloshin-A-I, Ostakhov-S-S.

Author affiliation

Kazakov, V.P., Voloshin, A.I., Ostakhov, S.S., Inst. of Org. Chem., Acad. of Sci., Ufa.

Abstract

Anomalous behavior of **chemiluminescence** (CL) of **adamantylidenadamantane-1,2-dioxetane** (Di) activated by chelate Eu (fod) branched quantum-chain processes of decomposition of dioxetane. The growth of the CL intensity with an increase in the Eu(III) concentration is well described by an exponential. On the basis of data on the kinetics of photophysical processes and the analysis of variations of spectral bands of Eu(III) **fluorescence** /sup 5/D/sub 1 /rightward arrow the dioxetane concentration, a conclusion is made that quenching of the complex with excitation in the /sup 5/D/sub 0/ state conserved leads to the decay of a hyperenergetic molecule of dioxetane in complex with Eu (III) and its excitation to the /sup 5/D/sub 1/ and.

Descriptors

CHEMILUMINESCENCE; FLUORESCENCE; ORGANIC-COMPOUNDS; RADIATION-QUENCHING; REACTION-KINETICS; SORPTION.

Classification codes

A8240T Chemiluminescence-and-chemical-laser-kinetics*;

A7860P Chemiluminescence-condensed-matter;

A3350D Molecular-fluorescence-and-phosphorescence-spectra;

A7855K Photoluminescence-in-organic-materials;

A8265M Sorption-and-accommodation-coefficients-surface-chemistry;

A8220 Chemical-kinetics.

Keywords

branched-quantum-chain-processes; 12-dioxetane; Eu(fod)/sub-3/; sorbent; anomalous-behavior; **chemiluminescence**; **adamantylidenadamantane-12-dioxetane**; chelate-Eu(fod)/sub-3/;

sorbent- surface; Eu(III)-concentration; kinetics; photophysical-processes; spectral-bands; **Eu(III)-fluorescence**; /sup-5/D/sub-1/rightward-arrow /sup-7/F/sub-j/; /sup-5/D/sub-0/rightward-arrow/sup-7/F/sub-j/; dioxetane-concentration; /sup-5/D/sub-1/-state-quenching; hyperenergetic-molecule.

Treatment codes

I Theoretical-or-mathematical;
X Experimental.

Language

English.

Publication type

Journal-paper.

Availability

SICI: 0030-400X(199711)83:5L.797:BQCP; 1-9.

CCCC: 0030-400X/96/835-0797\$10.00.

SICI of original-language version: 0030-4034(199711)83:5L.860; 1-7.

Publication year

1997.

Publication date

19971100.

Edition

1998002.

Copyright statement

Copyright 1998 IEE.

(c) 2007 The Institution of Engineering and Technology

Full text available at



SCIENCE @ DIRECT

USPTO Full Text Retrieval Options

open url

☒ document 52 of 65 [Order Document](#)

Inspec - 1898 to date (INZZ)

Accession number & update

0001923863 20051201.

Title

Optical activity due to deuterium substitution in **adamantanone** and **adamantanethione**. Barrier to inversion in the /sup 1/np π * excited state.

Source

Chemical Physics, {Chem-Phys-Netherlands}, 15 July 1982, vol. 69, no. 1-2, p. 19-26, 25 refs, CODEN: CMPHC2, ISSN: 0301-0104, Netherlands.

Author(s)

Schippers-P-H, Dekkers-H-P-J-M.

Author affiliation

Schippers, P.H., Dekkers, H.P.J.M., Dept. of Theoretical Organic Chem., Univ. of Leiden, Leiden, Netherlands.

Abstract

The circular dichroism (CD) and absorption were measured in the **(1S)-4,4-dideuterio-adamantan-2-one** (1) and **(1S)-4,4-dideuterio- adamantane-2-thione** (2). In addition the circular polarization in the **fluorescence** of 1 is reported. Since 1 and 2 owe their **optical** activity to a small perturbation of the C/sub 2v/ symmetry due to deuterium substitution, it is argued that the CD originates from transitions to totally symmetric upper state vibrational levels, whereas the absorption is induced by non-totally symmetric vibrations. The vibrational structure in the spectra is analysed in terms of the (thio) carbonyl stretching and out-of-plane bending modes. By comparison of the frequencies of the vibrations, active in CD and absorption, it is found that in the /sup 1/np π * state 1 has a double minimum potential in the out-of-plane bending mode, with an inversion barrier of ca. 1100 cm/sup -1/. The data for 2 are consistent with a much lower barrier to inversion.

Descriptors

~~MOLECULAR-ELECTRONIC-STATES~~; ~~MOLECULAR-VIBRATION~~; ~~OPTICAL-ROTATION~~;
~~ORGANIC-COMPOUNDS~~; ~~ROTATIONAL-ISOMERISM~~.

Classification codes

A3150 Excited-states-of-atoms-and-molecules*;
A3345 Magneto-optical-and-electro-optical-effects-in-molecules- birefringence-dichroism-and-optical-activity;
A3350D Molecular-fluorescence-and-phosphorescence-spectra;
A4210N Optical-polarization-in-homogeneous-media.

Keywords

optical-activity; D-substitution; isotope-effect; light-absorption; **fluorescence-polarisation**;
adamantanone; **adamantanethione**; circular- dichroism; **(1S)-44-dideuterio-adamantan-2-one**;
(1S)-44-dideuterio- **adamantane-2-thione**; circular-polarization; totally-symmetric-upper- state-
vibrational-levels; non-totally-symmetric-vibrations; vibrational-structure; carbonyl-stretching; out-of-
plane-bending-modes; double-minimum-potential; out-of-plane-bending-mode; inversion-barrier.

Treatment codes

X Experimental.

Language

English.

Publication type

Journal-paper.

Publication year

1982.

Publication date

19820715.

Edition

1982010.

Copyright statement

Copyright 1982 IEE.

(c) 2007 The Institution of Engineering and Technology

Full text
available
at

AMERICAN INSTITUTE OF PHYSICS



Electronic
Journals
Service

USPTO Full Text Retrieval Options

☒ **document 59 of 65** Order Document

Inspec - 1898 to date (INZZ)

Accession number & update

0000666475 20051201.

Title

Free radicals in an **adamantane** matrix. IX. **Optical** excitation and emission spectra of the benzyl radical and its monofluorinated derivatives.

Source

Journal of Chemical Physics, {J-Chem-Phys-USA}, 1 April 1974, vol. 60, no. 7, p. 2684-91, 29 refs, CODEN: JCPSA6, ISSN: 0021-9606, USA.

Author(s)

Lloyd-R-V, Wood-D-E.

Author affiliation





Lloyd, R.V., Wood, D.E., Univ. Connecticut, Storrs, CT, USA.

Abstract

For pt. XIII, see J. Am. Chem. Soc., vol.96, 659 (1974). Electronic emission- and excitation-spectra of the benzyl radical in an **adamantane** matrix resemble those found in other condensed phases, and are more readily obtained because of the ease with which operations using **adamantane** can be carried out. Corresponding spectra are obtained for the alpha-fluorobenzyl radical, and of the o-, m-, and p-fluorobenzyIs, some of these for the first time. Values for all substituted radicals (except the 3100 Å for p-fluorobenzyl) are shifted towards the red compared with the unsubstituted radical. Vibrational fine-

structure has been analysed for two bands of each radical.

Descriptors

 FREE-RADICALS;  MOLECULAR-VIBRATION;  ORGANIC-COMPOUNDS;  SPECTRA-OF-ORGANIC-MOLECULES-AND-SUBSTANCES.

Classification codes

A3320K Visible-molecular-spectra*;
A3520S Molecular-hyperfine-and-fine-structure-constants;
A7840 Visible-and-ultraviolet-spectra-condensed-matter.

Keywords

free-radicals; **adamantane-matrix**; emission-spectra; benzyl-radical; monofluorinated-derivatives;
optical-excitation-spectra; spectral-shift; vibrational-fine-structure.

Treatment codes

X Experimental.

Language

English.

Publication type

Journal-paper.

Publication year

1974.

Publication date

19740401.

Edition

1974007.

Copyright statement


Copyright 1974 IEE.

(c) 2007 The Institution of Engineering and Technology

previous
documents

next
documents

save

locally as: PDF document 

search strategy: do not include the search strategy 

order

copy to
Clipboard

Top - News & FAQs - Dialog

© 2007 Dialog